

## Medications to stop prior to your procedure

### **Please read instructions below carefully.**

Below is a list of medications the patient need to stop before the procedure. Please call with any questions or concerns regarding stopping these medications. **Please notify the office as soon as possible if you forget to stop any medications prior to your procedure.** Your procedure may or may not need to be rescheduled, depending on the circumstances.

**Please check with the prescribing physician of these medications and get permission from them to stop.** If you or the prescribing provider have any concerns regarding stopping these medications, they may call our office directly at **903-747-4737**.

**For diabetic patients:** Please check with your endocrinologist or primary care physician regarding your diabetic medications and how they should be taken while you prep for your procedure.

**The following medications need to be stopped *seven days prior (weight loss meds—14 days prior)*:**

Adipex	Clinoril	Naproxen
Advil	Clopidogrel	Naprosyn
Aggrenox	Coumadin	Orudis
Aleve	Ecotrin	Oruvail
Alka Seltzer	Effient	Persantin
Anaprox	Eliquis	<b>Phentermine</b>
Ansaid	Enoxaparin	<b>Phentarmine</b>
Apixaban	Excedrin w/ Aspirin	Plavix
Arthrotec	Feldene	Pletal
Aspirin 325 mg (81 mg is ok)	Heparin	Pradaxa
BC Powder	Ibuprofen	Prasugrel
Bextra	Indomethacin	Relafen
Brilinta	Ketorolac	Rivaroxaban
Cataflam/Cambia	Lodine	Ticlid
Celebrex	Lovenox	Ticagrelor
Celecoxib	Meloxicam	Toradol
Cilostazol	Mobic	Warfarin
	Motrin	Xarelto

Thank you for choosing UT Health Tyler. We look forward to seeing you soon.